

On the Standard Approach to Renormalization Group Improvement

F.A. Chishtie*, V. Elias†, R.B. Mann‡, D.G.C. McKeon§, T.G. Steele¶

February 2, 2008

UWO-TH-06/15

Abstract

Two approaches to renormalization-group improvement are examined: the substitution of the solutions of running couplings, masses and fields into perturbatively computed quantities is compared with the systematic sum of all the leading log (LL), next-to-leading log (NLL) *etc.* contributions to radiatively corrected processes, with n -loop expressions for the running quantities being responsible for summing $N^n LL$ contributions. A detailed comparison of these procedures is made in the context of the effective potential V in the 4-dimensional $O(4)$ massless $\lambda\phi^4$ model, showing the distinction between these procedures at two-loop order when considering the NLL contributions to the effective potential V .

The process of renormalization necessarily introduces an arbitrary renormalization scale μ into radiative processes in quantum field theory. Requiring that a process $\Gamma = \Gamma(\lambda, m, \phi, \mu)$ (with λ , m , ϕ being coupling, mass and external field) be independent of μ leads to the usual renormalization group (RG) equation.

$$\mu \frac{d\Gamma}{d\mu} = \left(\mu \frac{\partial}{\partial \mu} + \beta(\lambda) \frac{\partial}{\partial \lambda} + \gamma_m(\lambda) m \frac{\partial}{\partial m} + \gamma_\phi(\lambda) \phi \frac{\partial}{\partial \phi} \right) \Gamma = 0 \quad (1)$$

where

$$\beta = \mu \frac{d\lambda}{d\mu}, \quad \lambda(\mu_0) = \lambda_0, \quad (2)$$

$$\gamma_m = \frac{\mu}{m} \frac{dm}{d\mu}, \quad m(\mu_0) = m_0 \quad (3)$$

$$\gamma_\phi = \frac{\mu}{\phi} \frac{d\phi}{d\mu}, \quad \phi(\mu_0) = \phi_0 \quad (4)$$

The usual procedure [1] is to solve Eq. (1) by the replacement

$$\Gamma(\lambda, m, \phi, \mu) \rightarrow \Gamma(\bar{\lambda}(t), \bar{m}(t), \bar{\phi}(t), \mu) \quad (5)$$

*Department of Applied Mathematics, The University of Western Ontario, London, ON N6A 5B7, Canada

†Department of Applied Mathematics, The University of Western Ontario, London, ON N6A 5B7, Canada

‡Department of Physics, University of Waterloo, Waterloo, ON N2L 3G1, Canada

§Department of Applied Mathematics, The University of Western Ontario, London, ON N6A 5B7, Canada

¶Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, SK, S7N 5E2, Canada

by invoking the method of characteristics [2]. Here $\Gamma(\lambda, m, \phi, \mu)$ is a perturbatively computed approximation to the exact expression for Γ , and $\bar{\lambda}$, $\bar{m}$ and $\bar{\phi}$ are the solutions of the running versions Eqs. (2)–(4)

$$\frac{d\bar{\lambda}}{dt} = \beta(\bar{\lambda}(t)) \quad , \quad \bar{\lambda}(0) = \lambda_0 \quad (6)$$

$$\frac{1}{\bar{m}} \frac{d\bar{m}}{dt} = \gamma_m(\bar{\lambda}(t)) \quad , \quad \bar{m}(0) = m_0 \quad (7)$$

$$\frac{1}{\bar{\phi}} \frac{d\bar{\phi}}{dt} = \gamma_\phi(\bar{\lambda}(t)) \quad , \quad \bar{\phi}(0) = \phi_0 \quad (8)$$

where $t = \log \frac{A^2}{\mu^2}$ with A^2 typically (though not uniquely) specified by the energy scales of the process.

The incorporation of successively higher loop contributions to the functions β , γ_m and γ_ϕ into $\Gamma(\bar{\lambda}(t), \bar{m}(t), \bar{\phi}(t), \mu)$ represents a standard method of RG improvement. A detailed outline of this approach appears in Ch. 7 of Ref. [3], that has been applied, for example, to the contribution of the effective potential V in [4]. In this note we wish to examine this procedure in more detail, and illustrate how its application compares with a systematic sum of all the leading log (LL), next-to-leading log (NLL) *etc.* contributions. In particular, we look at the effective potential V in the $O(4)$ massless $\lambda\phi^4$ model¹ beyond one loop order in the NLL approximation.

The tree-level potential for this model is

$$V_{tree} = \pi^2 \phi^4 y \quad (9)$$

where $y \equiv \frac{\lambda}{4\pi^2}$ denotes the scalar couplant in the theory. At one loop order we have [5],²

$$\frac{d\bar{y}(t)}{dt} = 6\bar{y}(t)^2 \quad , \quad \bar{y}(0) = y \quad (10)$$

$$\frac{d\bar{\phi}(t)}{dt} = 0 \quad , \quad \bar{\phi}(0) = \phi \quad (11)$$

Replacing y and ϕ in Eq. (9) by the solutions to Eqs. (10–11)

$$\bar{y}(t) = y + 6y^2 t + O(t^2) \quad (12)$$

$$\bar{\phi}(t) = \phi + O(t) \quad . \quad (13)$$

results in

$$V_{tree} \rightarrow \pi^2 \bar{\phi}(t)^4 \bar{y}(t) = \pi^2 \phi^4 (y + 6y^2 t + \dots) \quad (14)$$

This expression agrees with the leading-logarithm (LL) terms in the explicit calculation of the one-loop potential [6, 7, 8, 9],

$$V_{1loop} = \pi^2 \phi^4 (y + 6y^2 t + T_{1,0} y^2 + \dots) \quad (15)$$

where $T_{1,0}$ denotes the non-logarithmic term whose value depends on the renormalization scheme.³ Thus the replacement of Eq. (14) does represent a systematic incorporation of the LL terms of the effective potential.

¹The $O(4)$ model is of particular interest as it represents the scalar field theory projection of the Standard Model.

²We have dropped the subscript “0” on y and ϕ for subsequent analysis.

³We have taken $t = L = \frac{1}{2} \log \left(\frac{\phi^2}{\mu^2} \right)$.

The general form of V is given by

$$V = \pi^2 \sum_{n=0}^{\infty} \sum_{m=0}^n y^{n+1} T_{n,m} t^m \phi^4 = \pi^2 \sum_{n=0}^{\infty} y^{n+1} S_n(yt) \phi^4, \quad T_{0,0} = 1 \quad (16)$$

where $S_n(\xi) = \sum_{m=0}^{\infty} T_{n+m,m} \xi^m$. In Refs. [10, 11] it is shown how Eq. (1) results in a series of ordinary differential equations that can be solved in turn for $S_0(\xi)$, $S_1(\xi)$, $\dots$ in terms of the functions $\beta(\lambda)$ and $\gamma_\phi(\lambda)$, provided that the boundary values $T_{n,0} = S_n(0)$ are known. The function $S_n(\lambda L)$ gives the $N^n LL$ contributions to V . In particular, $S_0 = \frac{1}{1-6yt}$ and $S_1 = \frac{4T_{1,0}-6yt+13\log(1-6yt)}{4(1-6yt)^2}$, so that to order t , we obtain

$$V_{LL} + V_{NLL} = \pi^2 \phi^4 (y S_0 + y^2 S_1) \approx \pi^2 \phi^4 [y + T_{1,0} y^2 + (6y^2 + 12T_{1,0} y^3 - 21y^3) t + O(t^2)] \quad (17)$$

which is indeed the NLL approximation to the explicit two-loop contributions to V

$$V_{1loop} + V_{2loop} = \pi^2 \phi^4 [y + T_{1,0} y^2 + T_{2,0} y^3 + (6y^2 + 12T_{1,0} y^3 - 21y^3) t + O(t^2)] \quad (18)$$

The quantity $T_{2,0}$, which is explicitly computed by a two loop calculation, is the initial value for the $NNLL$ summation $S_2(yt)$. As noted previously, we see from the LL terms in the above expression that the replacement of Eq. (14) does in fact generate the LL expression $V_{LL} = \pi^2 \phi^4 y S_0(yt)$. We now will check to see if a similar replacement of λ and ϕ by the two loop contribution to $\bar{\lambda}(t)$ and $\bar{\phi}(t)$ can generate the NLL results of either Eq. (17) or (18). The relevant RG equations at two-loop level are [14],

$$\frac{d\bar{y}(t)}{dt} = 6\bar{y}(t)^2 - \frac{39}{2}\bar{y}(t)^3 \quad (19)$$

$$\frac{d\bar{\phi}(t)}{dt} = -\frac{3}{8}\bar{y}(t)^2 \bar{\phi}(t) \quad (20)$$

whose solutions are

$$\bar{y}(t) = y + \left(6y^2 - \frac{39}{2}y^3\right) t + O(t^2) \quad (21)$$

$$\bar{\phi}(t) = \phi - \frac{3}{8}y^2 \phi t + O(t^2) \quad (22)$$

Replacing ϕ and y in V_{1loop} of Eq. (15) with the above results for $\bar{\phi}(t)$ and $\bar{y}(t)$ results in

$$V_{1loop} \rightarrow \pi^2 \phi^4 \left[y + T_{1,0} y^2 + \left(12y^2 - 21y^3 + 12T_{1,0} y^3 - \frac{81}{2}T_{1,0} y^4\right) t + O(t^2) \right] \quad (23)$$

which is distinct from both Eqs. (17) or (18) at LL level. Furthermore, substitution of equations (21) and (22) into the modified tree potential $\tilde{V}_{tree} = \pi^2 \phi^4 (y + T_{1,0} y^2)$ results in

$$\tilde{V}_{tree} \rightarrow \pi^2 \bar{\phi}(t)^4 [\bar{y}(t) + T_{1,0} \bar{y}(t)^2] \approx \pi^2 \phi^4 \left[y + T_{1,0} y^2 + \left(6y^2 + 12T_{1,0} y^3 - 21y^3 - \frac{81}{2}T_{1,0} y^4\right) t + O(t^2) \right] \quad (24)$$

This expression does agree with the NLL terms in (17) and (18), but it is not a systematic NLL expansion because is contaminated with $N^2 LL$ $y^4 t$ terms.

We thus see that simply replacing, in the one loop result, renormalized parameters by running parameters computed to two loop order, does not reproduce the systematic NLL approximation to (18) obtained either from explicit calculation of the two loop graphs or in (17) from carefully solving the RG equation for the functions S_0 or S_1 . In Section 7.6 of Ref. [3] the recursion relations that fix the form of the functions S_0 and S_1 are obtained from the RG equation and a comparison with the substitution of “running” RG group functions into perturbative results is suggested. It is this comparison that we have examined in detail above. One should not conclude from our analysis that the method of characteristics is inappropriate for extracting information from the renormalization group about physical processes. In Refs. [12, 13] ways in which this technique can be usefully applied are discussed.

It is worth briefly recapping the results of [12, 13] that are pertinent to the discussion above. We first illustrate the method of characteristics by considering the general equation

$$f(x, y) \frac{\partial A(x, y)}{\partial x} + g(x, y) \frac{\partial A(x, y)}{\partial y} + h(x, y) A(x, y) = 0 \quad (25)$$

where f , g and h are known functions. If $A_0(x, y)$ is a solution to Eq. (25), then it can be easily verified that

$$A(\bar{x}(\tau), \bar{y}(\tau), \tau) = A_0(\bar{x}(\tau), \bar{y}(\tau)) \exp \left(\int_0^\tau h(\bar{x}(\tau'), \bar{y}(\tau')) d\tau' \right) \quad (26)$$

is also a solution, where $\bar{x}(\tau)$ and $\bar{y}(\tau)$ are “characteristic functions” satisfying

$$\frac{d\bar{x}(\tau)}{d\tau} = f(\bar{x}(\tau), \bar{y}(\tau)) , \quad \bar{x}(0) = x \quad (27)$$

$$\frac{d\bar{y}(\tau)}{d\tau} = g(\bar{x}(\tau), \bar{y}(\tau)) , \quad \bar{y}(0) = y. \quad (28)$$

In fact, we can show from Eqs. (25,27,28) that

$$\frac{d}{d\tau} A(\bar{x}(\tau), \bar{y}(\tau), \tau) = 0, \quad (29)$$

For example, if $f(x, y) = x$, $g(x, y) = y^2$, $h(x, y) = 0$ and $A_0(x, y) = x e^{\frac{1}{y}}$, then

$$\bar{x}(\tau) = x e^\tau \quad (30)$$

$$\bar{y}(\tau) = y(1 - y\tau)^{-1} \quad (31)$$

and

$$A(\bar{x}(\tau), \bar{y}(\tau)) = A_0(x, y) . \quad (32)$$

However, if we only know the perturbative approximation to $A_0(x, y)$

$$A_0^{(1)}(x, y) = x \left(1 + \frac{1}{y} \right) \quad (33)$$

then

$$A_0^{(1)}(\bar{x}(\tau), \bar{y}(\tau)) = x e^\tau \left(1 + \frac{1 - y\tau}{y} \right) \quad (34)$$

which reduces to the exact solution $A_0(x, y)$ for one particular value of τ , namely $\frac{1}{y}$. Furthermore, if we extrapolate from Eq. (33) assume the form of $A_0(x, y)$ to be

$$A_0(x, y) = x \sum_{n=0}^{\infty} \alpha_n \left(\frac{1}{y} \right)^n \quad (35)$$

then the original differential equation [Eq.(25)] leads to the recursion relation

$$\alpha_n = (n + 1) \alpha_{n+1} \quad (36)$$

which shows that eq.(35) is equivalent to $x e^{\frac{1}{y}}$ if $\alpha_0 = 1$.

The points that arise in our discussion of this simple equation have direct analogues when one analyzes the RG equation given in Eq. (1). We first note that λ , m and ϕ in Eq. (1) are “running” functions of μ with their μ dependence dictated by Eqs. (2–8). In applying the method of characteristics to Eq. (1), we define “characteristic” functions $\bar{\lambda}(\tau)$, $\bar{m}(\tau)$, $\bar{\phi}(\tau)$ by the equations

$$\beta(\bar{\lambda}(\tau)) = \frac{d\bar{\lambda}(\tau)}{d\tau}, \quad \bar{\lambda}(0) = \lambda(\mu) \quad (37)$$

$$\gamma_m(\bar{\lambda}(\tau)) = \frac{1}{\bar{m}(\tau)} \frac{d\bar{m}(\tau)}{d\tau}, \quad \bar{m}(0) = m(\mu) \quad (38)$$

$$\gamma_\phi(\bar{\lambda}(\tau)) = \frac{1}{\bar{\phi}(\tau)} \frac{d\bar{\phi}(\tau)}{d\tau}, \quad \bar{\phi}(0) = \phi(\mu). \quad (39)$$

In Eqs. (37–39), we see that the running functions act as boundary values for the characteristic functions, while the form of Eqs. (6–8) is identical to that of Eqs. (37–39).

Just as substitution of the characteristic functions of Eqs. (30,31) into the perturbative solution (33) reduces to the exact solution, so the substitution of characteristic functions satisfying Eqs. (37–39) into perturbatively computed results for V can reproduce exact solutions for V in terms of the functions $S_0, S_1, \dots$ when the characteristic parameter τ takes on a particular value. This is illustrated in the context of massless scalar electrodynamics in Section 7 of Ref. [13].

We also note that determining an exact solution to Eq. (25) through substitution of Eq. (35) into Eq. (25) and thereby obtaining the recursion relation of Eq. (36) is directly analogous to the way in which the functions S_0, S_1 etc. are generated in Refs. [10, 11].

Acknowledgements

One of the authors, V. Elias, has recently passed away; the other authors would particularly like to acknowledge their indebtedness to him, both in this work and in many other collaborative efforts with him. NSERC provided financial support.

References

- [1] S. Coleman, “Aspects of Symmetry”, Cambridge University Press, Cambridge (1985)
- [2] A. Peterman, Phys. Rep. C53 (1979) 157.
- [3] J.C. Collins, “Renormalization”, Cambridge University Press, Cambridge (1984).
- [4] J. A. Casas, J. R. Espinosa, and M. Quiros, Phys. Lett. B342 (1995) 171.
- [5] J. C. Collins, Phys. Rev. D10 (1974) 1213.
- [6] S. Coleman and E. Weinberg, Phys. Rev. D7 (1973) 1888.
- [7] S. Weinberg, Phys. Rev. D7 (1973) 2887.
- [8] R. Jackiw, Phys. Rev. D9 (1974) 1686.
- [9] A. Salam and J. Strathdee, Phys. Rev. D9 (1974) 1129.
- [10] B. Kastening, Phys. Lett. B283 (1992) 287.
- [11] V. Elias, R.B. Mann, D.G.C. McKeon and T.G. Steele, Nucl. Phys. B 678 (2004) 147; (Erratum) Nucl. Phys. B 703 (2004) 413.
- [12] V. Elias, D.G.C. McKeon, Int. J. Mod. Phys. A, 18 (2003) 3417.
- [13] V. Elias, D.G.C. McKeon and T.N. Sherry, Int. J. Mod. Phys. A, 20 (2005) 1065.
- [14] H. Kleinert, J. Neu, N. Schulte-Frohlinde, K.G. Chetyrkin, S.A. Larin, Phys. Lett. B 272 (1991) 39; (Erratum) Phys. Lett. B 319 (1993)